

Building portfolios of stocks in the São Paulo Stock Exchange using Random Matrix Theory

Leonidas Sandoval Junior ^{*}
 Adriana Bruscato [†]
 Maria Kelly Venezuela [‡]

Insper, Institute of Education and Research

January 4, 2012

Abstract

Using Random Matrix Theory, we build a covariance matrix between stocks of the BM&F-Bovespa (Bolsa de Valores, Mercadorias e Futuros de São Paulo) which is cleaned of some of the noise due to the complex interactions between the many stocks and the finiteness of available data, and use a regression model in order to remove the market effect due to the common movement of all stocks. These two procedures are then used in order to build portfolios of stocks based on Markovitz's theory, trying to build better predictions of future risk based on past data. This is done for years of both low and high volatility of the Brazilian market, from 2004 to 2010.

1 Introduction

Modern portfolio theory is largely based on Markovitz's ideas [1]-[3], where a portfolio of various equities is built on the principle of minimizing risk given an expected return. Risk is assessed as the volatility of each stock that comprises the portfolio, as well as their covariance. Preference is given to stocks that have negative or low covariance between each other, what leads to diversification of the equities held in one particular portfolio.

Both volatility and covariance are integrated into the covariance matrix, which is built using the stock returns of past data. This is used in order to predict the risk of a portfolio, and it is usually different from the realized risk of the same portfolio.

Three problems arise from this approach. The first one is that past data reflect the market as it was, and not as it will be. So, the theory assumes the hypothesis that future events shall mimic past events, what is usually not true, for it doesn't incorporate the release of news, or the current mood of the market. There is not much one can do about it, but in order to minimize effects of events that might change the behavior of a market, one cannot use past data that is too old.

This leads us to the second problem, which is the noise associated with past data that arises purely from the fact that the available data is finite. Since one cannot go back in time indefinitely, and even if one could, it wouldn't be advisable given the discussion in the preceding paragraph, there is only a limited amount of data (in our case, price quotations) from which to build a covariance matrix. The problem gets even more severe if we think that an efficient portfolio should be built from many and diverse equities. A third source of noise comes from the complex interactions between the many elements of a stock market: traders, news, foreign markets, and the very prices of stocks interact in order to guide the price of a stock. Those interactions are usually too complex to be accommodated by any economic model.

So, all this noise is incorporated into the covariance matrix that is used in order to try to forecast the risk of a particular portfolio, and if one can clean that matrix from some of that noise, one is then able to make

^{*}E-mail: LeonidasSJ@insper.edu.br (corresponding author)

[†]E-mail: AdrianaB@insper.edu.br

[‡]E-mail: MariaKV@insper.edu.br

better predictions for risk. Many studies have been made on the influence of noise and other factors on the covariance matrix in the building of portfolios [4]-[9]. Most of the approaches for solving them involve the reduction of the dimensionality of the covariance matrix by introducing some structure into it, obtained by principal component analysis, separation of stocks into economic sectors, among other means (see [10] and [11] for two of these approaches).

A technique first developed for the study of the nuclei of the atoms of heavier elements, called Random Matrix Theory [12], compares the eigenvalues of a correlation matrix with those of a correlation matrix built from a purely random matrix. From such a comparison, one may then discern elements which are clearly not random, and study them separately. Such technique has been applied to a number of complex systems, and, particularly, to financial markets. Of the many results that were obtained, the building of portfolios that resemble more closely the realized risk of the future market, based on past data, is one of them [13] [14], and it has been successfully applied to stocks [15] [16] and mutual funds [17].

Besides the cleaning of the correlation matrix, we used a regression model in order to remove the market effect on the asset returns. This procedure makes it possible that the correlation matrix be estimated with greater precision, for there is just a part of the dependence which is due to the assets, what generates more reliable forecasts for the risk of a portfolio.

The contribution of this article is the use of a method which is capable of ameliorating the risk forecasts of a portfolio built with assets of the Brazilian stock market, based on past data. This method involves three steps: (1) the removal of the market effect of the assets; (2) the cleaning of the correlation matrix, which encodes the structure of the dependence of the assets being considered, based on Random Matrix Theory, and (3) the construction of portfolios using Markowitz's theory and the cleaned correlation matrix. In this article, we calculate portfolios of stocks with and without the removal of the market effect so as to compare both results.

In order to analyze the suitability of the proposed method, we shall use the daily returns of stocks of the BM&F-Bovespa with 100% liquidity in which pairs of years ranging from 2004 to 2010. For each year being analyzed, we build a portfolio using data from the previous year in order to make a forecast of the risk for the a determined year, and that forecasted risk is then compared with the realized risk in that year. As data used in this article include periods of both low and high volatility in the BM&F-Bovespa, in particular the data collected during the Subprime Mortgage Crisis of 2007 and 2008, we are able to study how this technique of cleaning the correlation matrix applies to times of high volatility.

The article is organized in the following way: Section 2 introduces the basic concepts of Random Matrix Theory, which are then applied to the building of portfolios (according to Markowitz) with and without cleaning the correlation matrix for short selling allowed, as well as the regression for the removal of the market effect. Section 3 shows the results of portfolios forecasts using data from 2004 to 2010 and compares them to the realized risks. The article ends with a conclusion and comments on years of high volatility.

2 Methodology

In this section, we briefly describe the method proposed for the construction of portfolios by cleaning the correlation matrix and removing the market effect, aiming at a better forecasting of risk based on the previous behaviors of the assets. We use the year 2004 as an example of the application of such method in this section.

2.1 Random matrix theory

Random matrix theory had its origins in 1953, in the work of the German physicist Eugene Wigner [18] [19]. He was studying the energy levels of complex atomic nuclei, such as uranium, and had no means of calculating the distance between those levels. He then assumed that those distances between energy levels should be similar to the ones obtained from a random matrix which expressed the connections between the many energy levels. Surprisingly, he could then be able to make sensible predictions about how the energy levels related to one another.

The theory was later developed, with many and surprising results arising. Today, random matrix theory is applied to quantum physics, nanotechnology, quantum gravity, the study of the structure of crystals, and may have applications in ecology, linguistics, and many other fields where a huge amount of apparently unrelated

information may be understood as being somehow connected. The theory has also been applied to finance in a series of works dealing with the correlation matrices of stock prices, and also with risk management in portfolios [20]-[23] (for a recent review on the subject, see [24]).

The first result of the theory that we shall mention is that, given an $L \times N$ matrix with random numbers built on a Gaussian distribution with average zero and standard deviation σ , then, in the limit $L \rightarrow \infty$ and $N \rightarrow \infty$ such that $Q = L/N$ remains finite and greater than 1, the eigenvalues λ of such a matrix will have the following probability density function, called a Marčenko-Pastur distribution [25]:

$$\rho(\lambda) = \frac{Q}{2\pi\sigma^2} \frac{\sqrt{(\lambda_+ - \lambda)(\lambda - \lambda_-)}}{\lambda}, \quad (1)$$

where

$$\lambda_- = \sigma^2 \left(1 + \frac{1}{Q} - 2\sqrt{\frac{1}{Q}} \right), \quad \lambda_+ = \sigma^2 \left(1 + \frac{1}{Q} + 2\sqrt{\frac{1}{Q}} \right), \quad (2)$$

and λ is restricted to the interval $[\lambda_-, \lambda_+]$.

Figure 1 shows some of these distributions for diverse values of Q and σ .




Fig. 1. Marčenko-Pastur distribution for fixed values of Q and σ .

Since the distribution (1) is only valid for the limit $L \rightarrow \infty$ and $N \rightarrow \infty$, finite distributions will present differences from this behavior. Another source of deviations is the fact that financial time series are better described by non-Gaussian distributions, such as t Student or Tsallis distribution.

In Figure 2, we compare the theoretical distribution for $Q = 10$ and $\sigma = 1$ to distributions of the eigenvalues of three correlation matrices generated from finite $L \times N$ matrices such that $Q = L/M = 10$, and the elements of the matrices are random numbers with mean zero and standard deviation 1.




Fig. 2. Histogram of eigenvalues for generated correlation matrix and Marčenko-Pastur theoretical distribution (solid line) $Q = 10$ and $\sigma = 1$.

Consequently, real data will deviate from the theoretic probability distribution. Nevertheless, the theoretical result may serve as a parameter to the results obtained experimentally.

We shall now explain why random matrix theory is useful for portfolio building, starting by explaining how it can be used in order to clean the correlation matrix of some of its noise. In order to do that we shall consider the data concerning the year 2004, the first of the years considered here in our study. For this period we chose stocks of the Bovespa (by then not yet joined with the BM&F) which were negotiated every trading day during the years 2004 and 2005 (2004 will be the past data that will be used to predict the risk in 2005). Those stocks are listed in Appendix A, totalizing 61 stocks.

For each stock, we calculated the returns, more precisely the log-returns, given by

$$R_t = \ln(P_t) - \ln(P_{t-1}) \approx \frac{P_t - P_{t-1}}{P_t}, \quad (3)$$

where P_t are the closing prices of one stock at the trading day t . Those returns were then normalized in order to obtain zero mean and standard deviation one by using the formula

$$X_t = \frac{R_t - \mu_R}{\sigma_R}, \quad (4)$$

where μ_R is the average of the time series used, and σ_R is its standard deviation. This is done in order to best compare the resulting correlation matrix with the theoretical one, which is a random matrix with zero mean and standard deviation σ chosen to be equal to one. The correlation matrix (a 61×61 matrix) between the variables X_t for the year 2004 was then calculated.

The distribution density of the eigenvalues of the correlation matrix thus obtained is shown in Figure 3 (left picture). Also, the eigenvalues are plotted in order of magnitude in right picture of Figure 3. The shaded area indicates the region predicted by theory for the data related to a purely random behavior of the normalized returns.




Fig. 3. Left: histogram of eigenvalues for the correlation matrix of 62 stocks in 2004 and Marčenko-Pastur theoretical distribution (solid line). Right: eigenvalues for the correlation matrix of 62 stocks in 2004 and purely random region.

We have $L = 248$ days of data for each of the $N = 61$ stocks, so that $Q = 248/61 \approx 4,06$. The probability distribution function for a random matrix with $L \rightarrow \infty$ and $N \rightarrow \infty$ with $Q \approx 4,06$ is also plotted in Figure 3,

so that we may compare the result of pure noise with the one obtained for our data. The minimum (λ_-) and maximum (λ_+) values of the probability distribution function are given by

$$\lambda_- = 0.254 \quad \text{and} \quad \lambda_+ = 2.238 .$$

The first striking feature is that the largest eigenvalue is more than ten times larger than the maximum value predicted for a purely random correlation matrix. About 72% of the eigenvalues fall within the shaded region associated with pure noise, 15 of them fall below this region, and another one is above it.

The eigenvectors e_1 and e_2 for the two largest eigenvalues, $\lambda_1 = 23.505$, and $\lambda_2 = 2.540$, are represented in Figure 4 (first two graphics). The white bars represent positive values and the gray bars represent negative ones.

The distribution of individual values of eigenvector e_1 is very similar for all the stocks considered, showing that all stocks contribute to this mode, which is considered “the market mode”. For eigenvector e_2 , one can see the prevalence of some stocks over others. In comparison, eigenvectors corresponding to the shaded region (Wishart region) do not show any preference for particular stocks.

The third and fourth graphs of Figure 4 show the eigenvectors distributions for the eigenvalues of two eigenvectors that are inside the region considered as noise, $\lambda_{18} = 0.853$, and $\lambda_{37} = 0.393$. Note that there are no structures of stocks clearly defined.




Fig. 4. Eigenvectors of some fixed eigenvalues: λ_1 , λ_2 (largest), λ_{18} , λ_{37} (noise region), and λ_{60} , λ_{61} (lowest eigenvalues).

We also show the eigenvectors corresponding to the two lowest eigenvalues of the correlation matrix, $\lambda_{60} = 0,046$ and $\lambda_{62} = 0,039$ (last two graphs in Figure 4). These eigenvectors corresponding to low eigenvalues represent “portfolios” of low risk, in opposition to the eigenvectors of the largest two eigenvectors, which represent the oscillations of the market and the common behavior of a cluster of stocks that behave in a similar way. Eigenvector e_{61} represents a portfolio where the investor buys PETR4 and short-sells PETR3, which are stocks belonging to the same company, Petrobras, and buys ELET3 and short-sells ELET6, which are belong also to the same company, Eletrobras, and eigenvector e_{60} represents a portfolio where the investor buys VALE3 and ELET6 and short-sells in VALE5 and ELET3, which again are two pairs of stocks of the same companies, and also buys PETR3 and short-sells PETR4.

Figure 5 shows the daily log-returns of portfolio P_1 built with eigenvalue e_1 , plotted against the log-returns of the Ibovespa, which is an index that describes the general behavior of the São Paulo Stock Exchange. The correlation between the two vectors is 0.9865, which is a very strong indication that the portfolio P_1 corresponds to a combination of stocks that behave much like the market, although with a much larger volatility: the standard deviation of the returns of P_1 is 12.51%, and the standard deviation for the Ibovespa is 1.80%.

The situation changes if we consider a portfolio built with eigenvector e_{37} , which corresponds to the noisy part of the eigenvalue spectrum: the correlation between this portfolio, P_{37} , and the Ibovespa is 0.1824, and it has a standard deviation 1.72%, very close to the standard deviation of the Ibovespa. For the portfolio P_{61} , built with eigenvector e_{61} , which corresponds to the lowest eigenvalue, the correlation with the Ibovespa is 0.0932, and its standard deviation is 0.44%. This portfolio presents the lowest correlation with the Ibovespa.




Fig. 5. Scatter plot of the portfolio built with e_1 and Ibovespa returns.

2.2 Building portfolios using Markowitz’s Theory

In this section, we shall build portfolios using the $N = 61$ stocks we are considering based on the correlation matrix of their returns in the year 2004. According to the usual portfolio theory, we can obtain w , the vector of weights of the portfolio due to each stock, by fixing the portfolio return (RE) and minimizing the risk (RI) of the portfolio.

The return of the portfolio is given by

$$RE = w^T R , \quad (5)$$

where R is the vector of average returns of each stock in 2004.

The risk is defined by the variance of the portfolio

$$RI = w^T \Sigma_R w , \quad (6)$$

where Σ_R is the estimated covariance matrix of the N stocks

The risk is then minimized with the constraint that the sum of all weights in the portfolio should be equal to one,

$$\sum_{i=1}^N w_i = 1 . \quad (7)$$

One can do that for several values of the average return, leaving the coordinates of w free to assume negative values, as well as positive ones, so that short selling is allowed. In Finance, this is not always possible, or sometimes it is limited, and so we shall consider the case of no short selling later on.

In order to build a portfolio, the covariance matrix of a period of time (usually some months) prior to the period of investment is used together with a forecast of the expected returns. Those returns, which are unknown, may be approximated by many means, with relative degrees of success. There is a vast literature on the forecasting of returns [2], but this does not concern us in our study of how to improve the prediction of risk. So, in order to restrict ourselves to the analysis of the correlation matrix, we shall consider that our prediction of returns is the best possible one, which is a perfect forecast of returns. Of course, if we had a perfect forecast of returns, and we knew it was a perfect forecast, we wouldn't need to make any portfolio analysis. We use here the perfect forecast of returns in order to compare different ways of calculating risk in a way that is independent of the way one tries to forecast returns.

So, we first use the covariance matrix from 2004, together with the average returns of 2005 (perfect forecast of returns), in order to build minimum risk portfolios for 2005. Doing so, we build an *efficient frontier*, which is a curve whose coordinates are the minimum risk for a given return. We also use the data from 2005, which means perfect forecasts of risk and return, in order to build an efficient frontier for the realized risk.

In Figure 6, we have the predicted return and risk of portfolios using 2004 covariances of stocks (dashed line) and the realized return and risk using data from 2005 (solid line). Note that, for a given return, the predicted risk is smaller than the realized risk. This may lead to a false perception of how risky an investment truly is, and may cause wrong decisions by the manager of the portfolio. The agreement of the curves (predicted and realized risk) can be measured by

$$AG = \frac{1}{n} \sum_{i=1}^n \frac{RI_i^{real} - RI_i^{pred}}{RI_i^{pred}} , \quad (8)$$

where RI_i^{real} is the realized risk and RI_i^{pred} is the predicted risk, both for $i = 1, \dots, n$ values of fixed returns. In our case, this number is $AG = -0.176$ ($n = 100$), which means that the predicted risk is, in average, 18% smaller than the realized risk.




Fig. 6. Realized (solid line) and predicted (dashed line) returns and risks of portfolios for 2005.

2.3 Building portfolios with a cleaned correlation matrix

The situation may be improved by trying to remove some of the noise of the correlation matrices of the returns of 2004 and 2005. One way this can be done is by building a diagonal matrix D where the elements of the diagonal are the eigenvalues of the original correlation matrix, but now with all eigenvalues corresponding to noise (those between λ_- and λ_+) replaced by their average [13]-[17]. In our present case, this average is

$\bar{\lambda} = 0.748$ for the eigenvalues based on data from 2004 and $\bar{\lambda} = 0.790$ for the eigenvalues based on data from 2005. The cleaned correlation matrix is then built using the formula

$$\mathbb{C}_{\text{clean}} = PDP^{-1} , \quad (9)$$

where P are matrices whose columns are the eigenvectors of the original correlation matrix.

Calculating now the efficient frontier built with the covariance matrix obtained from the cleaned correlation matrix of 2004, together with the average returns of 2005 (perfect forecast of returns), dashed line, and comparing with the real curve calculated with the covariance matrix obtained from the cleaned correlation matrix of 2005, solid line, we obtain the results represented in Figure 7.

The ratio between predicted and realized risk has now gone from $AG = -0.176$ to $AG = -0.102$, which means that the predicted risk is, in average, 10% smaller than the realized risk. This is an improvement on the previous result and shows how the cleaning of the correlation matrix may help building portfolios which account best for the realized risk based on previous data.




Fig. 7. Realized (solid line) and predicted (dashed line) returns and risks of portfolios for 2005 based on the cleaned correlation matrices.

2.4 Systemic risk

One of the stylized facts that remains, concerning the returns of assets, is the existence of volatility clustering. Generally, those volatility clusterings occur due to patterns of the stock market, and they make it difficult to forecast the risk of portfolios, since they show a structure of dependence between the assets and the market, and not solely the dependence between assets. As an example, the prediction for 2008 using data from 2007 grossly underestimates the risk of 2008, for 2007 was a year with relatively low volatility and 2008 saw the height of the USA Subprime Mortgage Crisis. Similarly, risk prediction for 2009 using data from 2008 overestimates the risk for 2009. Figure 8a shows the volatility of the Ibovespa (the index for the São Paulo Stock Exchange) over the years considered in this article. The volatility is calculated as the absolute value of the log returns of the index. It clearly shows regions with high and low volatilities.

The most common way to remove this so called systemic risk is to use a single index model, where all log returns R_t are written in terms of a first degree function of a market index I_t , as, for example, the Ibovespa, plus a residue E_t :

$$R_t = a + bI_t + E_t . \quad (10)$$

The coefficients a and b are determined for each equity using simple linear regression and the residue E_t is defined by equation 10.

As an alternative to the use of the Ibovespa as the single index, one may use the index obtained by the log returns of the portfolio of stocks that may be built using the eigenvector corresponding to the highest eigenvalue of the correlation matrix of those same stocks. As we have shown for the data concerning the year 2004, both this index and the Ibovespa are very highly correlated, so the results shouldn't be substantially altered by using any of those two indices.

Figures 8b and 8c show, respectively, the volatility of the log returns and of the residuals of PETR4, stocks from Petrobras, a gas and oil company that is one of the largest assets of the BM&F-Bovespa in terms

of negotiated volume. Note that the volatility of returns is less prone to effects due to market fluctuations, although those effects are still present in its variability. This remaining nonstationarity shall have its effects on the results in the next section.




Fig. 8. Volatility of Ibovespa (a), PETR4 (b) and the residuals of PETR4 (c) from 2004 to 2010.

So, in what follows, we calculate the residuals for all stocks being considered for each pair of years using the Ibovespa as the single index. We then proceed to building portfolios using the correlation matrices between those residuals and also the cleaned correlation matrices.

3 Results

In this section, we shall describe the results obtained with the procedure presented in Section 2 for each of the years considered in this article. We shall make a comparison of the predicted risk and the realized risk for the years 2005 to 2010, using graphics and calculating AG .

Table 1 presents the results for the measure AG calculated for this method with and without the cleaning of the correlation matrix and with and without the regression for the removal of the market effect, as well as the volatility of the Ibovespa.

Previous-Predicted	Without Cleaning		With Cleaning		Ibovespa Volatility
	No Regression	Regression	No Regression	Regression	
2004 – 2005	−0,176	−0,105	−0.102	0,021	1,57%
2005 – 2006	−0.212	−0,079	−0,203	0,064	1,53%
2006 – 2007	−0.071	0,018	−0.056	−0,101	1,73%
2007 – 2008	−1.011	−0,220	−0.841	−0,283	3,32%
2008 – 2009	0.259	0,290	0.169	0,263	1,93%
2009 – 2010	−0,056	−0,218	−0,116	−0,148	1,28%

Table 1. Agreement measure of the curves (AG) and the volatility of the Ibovespa.

One could notice that in Table 1 the majority of results for forecasts were better with the use of regression in order to eliminate the effect of the market movements and with the cleaning of the correlation matrix. Moreover, in every year in which there was a volatility drop from the previous year to the forecasted year (2005, 2006, 2009 and 2010), the cleaning of the correlation matrix showed better results, in other words, when volatility drops with respect to the previous year, it is best to use the cleaning of the correlation matrix, and when volatility grows with respect to the previous year, it is best not to use the cleaning of the correlation matrix. So, an investor may judge which would be the best method based on his anticipations in terms of the volatility of the market or use both procedures, with and without the cleaning the correlation matrix, in order to decide his or her strategy.

Figure 9 contains the graphs of the return and realized risk (solid line), and predicted (dashed line) with the original correlation matrix (left plots) and the cleaned correlation matrix (right plots), always using regression in order to remove the market effect. Based on these graphics, one may notice that the cleaning procedure was best for years of low volatility of the Ibovespa.

The year 2005 was relatively quiet for the Bovespa (by then, it hadn't yet merged with the BM&F). So, there was low volatility for 2005, and the predictions of risk are better using the residuals of data concerning this year. For this period, we chose stocks of the Bovespa which were negotiated every trading day during the years 2004 and 2005. Those stocks are listed in Appendix A, totalizing 61 stocks. For the correlation matrix of 2004 and the returns of 2005, we obtain $AG = -0.105$, which means that the predicted risk is 10% smaller than the realized risk. For the cleaned correlation matrices, $AG = 0.021$, which means that the predicted risk is 2% larger than the realized risk.

The year of 2006 was also a low volatility year for the Bovespa. For this period, we chose stocks of the Bovespa which were negotiated every trading day during the years 2005 and 2006. Those stocks are listed in Appendix A, totalizing 72 stocks. The predicted risk is 8% smaller than the realized risk for the original correlation matrix, and for the cleaned correlation matrix, the predicted risk is 6% larger than the realized risk.

Although 2006 was a year of low volatility, 2007 saw the beginning of the USA Subprime Mortgage Crisis, so there is a strong difference in volatility between those two years even when removing the market effect. For this period we chose stocks of the Bovespa which were negotiated every trading day during the years 2006 and 2007, listed in Appendix A, totalizing 86 stocks. For the original correlation matrix the predicted risk is 2% larger than the realized risk and for the cleaned correlation matrix, the predicted risk is 10% smaller than the realized risk.



The year 2008 was the most turbulent year in Brazil and in stock exchanges worldwide since the Black Monday crisis of 1987. The USA Subprime Mortgage Crisis spread to become a crisis of trust in financial institutions and to a worldwide credit crisis. For this period, we chose stocks of the Bovespa (BM&F-Bovespa from 2008 onwards) which were negotiated every trading day during the years 2007 and 2008, also listed in Appendix A, totalizing 105 stocks. For the original correlation matrix, the predicted risk is 22% smaller than the realized risk, and for the cleaned correlation matrix, the predicted risk is 28% smaller than the realized risk. From all procedures, this was the year that resulted in the worst forecasts.

Volatility remained high for 2009, although in Brazil the international economic crisis did not have the same strength as in other countries. So, the BM&F-Bovespa had a quieter period than some other stock exchanges, although the movement in and out of foreign investment remained high. For this period, we chose stocks of the



Bovespa which were negotiated every trading day during the years 2008 and 2009. Those stocks are listed in Appendix A, totaling 148 stocks. For the original correlation matrix the predicted risk is 29% larger than the realized risk, and for the cleaned correlation matrices the predicted risk is 26% larger than the realized risk.

Volatility remained higher than normal for the year 2010, and apprehension due to a succession of international financial crises made the market unstable, but less than in the previous years. For this period we chose stocks of the Bovespa which were negotiated every trading day during the years 2009 and 2010, listed in Appendix A, totaling 153 stocks. For the original correlation matrix the predicted risk is 22% larger than the realized risk, and for the cleaned correlation matrix the predicted risk is 15% larger than the realized risk.



After the crisis of 2008, the forecasts became poorer when compared with the ones of the previous years, indicating that the highest the change in volatility in subsequent years, the worst are the forecasts obtained for portfolios.





(a) 2005





(b) 2006





(c) 2007



(d) 2008



(e) 2009



(f) 2010

Fig. 9. Realized (solid lines) and predicted (dashed lines) returns and risks of portfolios using the original correlation matrices (left plots) and the cleaned correlation matrices (right plots) from 2005 to 2010.

As mentioned before, short selling is not usually freely allowed in financial transactions, mainly due to the increase in risks it might bring to a portfolio. So, we also did the calculations for no short selling, which implies in the addition of a constraint that specifies that all values in the weight vector w must be positive or zero. The results are summarized in Table 2, and they show that the conclusions to be derived from the case with

no short selling allowed are basically the same as the ones from the case where short selling is allowed: there is usually a significant better result when removing the market effect through a regression and, in years of a small variation of volatility, the cleaning of the correlation matrices is a better method for forecasting risk, and it fails for the forecasting of risk for years of high volatility.

Previous-Predicted	Without Cleaning		With Cleaning		Ibovespa Volatility
	No Regression	Regression	No Regression	Regression	
2004 – 2005	0.057	−0,066	0.069	0,001	1,57%
2005 – 2006	−0.128	0.041	−0,104	0,074	1,53%
2006 – 2007	−0.263	0,133	−0.261	−0,178	1,73%
2007 – 2008	−0.462	−0,564	−0.467	−0,606	3,32%
2008 – 2009	0.506	0,322	0.514	0,337	1,93%
2009 – 2010	−0,149	0,207	−0,136	0,176	1,28%

Table 2. Agreement measure of the curves (AG) and the volatility of the Ibovespa without the use of short selling.

4 Conclusion

In this article, we used two techniques in order to clean the covariance matrix used in the building of portfolios using Markovitz’s theory. The first technique is the use of Random Matrix Theory in order to clean the correlation matrix built from the time series data of stocks in the year prior to the year for which the portfolio is to be built. The second technique is to use a regression model in the removal of the market effect due to the common movement of all stocks. These are used in order to forecast the risk of a portfolio in a particular year using data from its previous year with better precision. The data were the time series returns of the 100% liquid assets of the BM&F-Bovespa covering the years from 2004 to 2010. The aim was to combine these two methods in different configurations, and to compare these results in order to obtain the best risk forecasts for portfolios.

Based on a measure of the agreement (AG) between the forecasted and the realized risks, we conclude that the forecasted risk is closer to the realized risk, depending on the volatility of the forecasted year being smaller or larger than the one of the year used for the forecast. In the case there is a drop in volatility from one year to the other, the results of the agreement measure were best when we built a portfolio using the cleaning of the correlation matrices. In the case there is a raise in volatility, it is best to use the original correlation matrix. This result is valid when the market effect is both removed and maintained by a regression.

The better performance of the method of cleaning the correlation matrices for the building of portfolios occurs because it eliminates the volatility (noise due to the complex interactions between the many stocks) which is useless for the forecasting of risk for a subsequent year of lower volatility. We noticed the advantage, in slightly more than half the cases, in making the forecast of risk having the market effect removed, but without the presence of a clear pattern. Last, in the year of the greatest crisis (2008), the use of regression was particularly better in terms of previsibility.

So, the use of regression methods in the removal of market effects is usually advisable, but the use of Random Matrix Theory in the removal of noise from the correlation matrices tend to fail in the forecasting for years of high volatility, which are precisely the occasions in which a reliable risk forecast is most needed.

References

- [1] H.M. Markowitz, *Portfolio Selection*, The Journal of Finance 7 (1) (1952), 77-91.
- [2] E.J. Elton, M.J. Gruber, S.J. Brown, and W. Goetzmann, *Modern Portfolio Theory and Investment Analysis*, Eighth Edition , (2009) Wiley.
- [3] Z. Bodie, A. Kane, and A.J. Marcus, *Investments*, Eighth Edition , (2009) McGraww-Hill/Irwin.

- [4] G.M. Frankfurter, H.E. Phillips, and J.P. Seagle, *Portfolio Selection: The Effects of Uncertain Means, Variances, and Covariances*, J. Fin. Quant. Anal. **6** (1971) 1251-1262.
- [5] G.M. Frankfurter, H.E. Phillips, and J.P. Seagle, *Estimation Risk in the Portfolio Selection Model: A Comment*, J. Fin. Quant. Anal. **7** (1972) 1423-1424.
- [6] J. P. Dickinson, *The Reliability of Estimation Procedures in Portfolio Analysis*, J. Fin. Quant. Anal. **9** (1974) 447-462.
- [7] J.D. Jobson and B.M. Korkie, *Estimation for Markowitz Efficient Portfolios*, J. Am. Stat. Assoc. **75** (1980) 544-554.
- [8] R.O. Michaud, *The Markowitz Optimization Enigma: Is 'Optimized' Optimal?*, Fin. Anal. Journal **45** (1989) 31-42.
- [9] V. Chopra and W.T. Ziemba, *The Effect of Errors in Mean and Co-Variance Estimates on Optimal Portfolio Choice*, J. Port. Management (1993) 6-11.
- [10] P. Jorion, *Bayes-Stein Estimation for Portfolio Analysis*, J. Fin. Quant. Anal. **21** (1986) 279-292.
- [11] V. DeMiguel, L. Garlappi, F.J. Nogales, and R. Uppal, *A Generalized Approach to Portfolio Optimization: Improving Performance by Constraining Portfolio Norms*, Management Science **55** (2009) 782-812.
- [12] M. L. Mehta, *Random Matrices*, (2004) Academic Press.
- [13] L. Laloux, P. Cizeau, J.-P. Bouchaud, and M. Potters, *Random Matrix Theory and Financial Correlations*, Int. J. Theor. Appl. Finance. **3** (2000) 391-397.
- [14] B. Rosenow, V. Plerou, P. Gopikrishnan, and H.E. Stanley, *Portfolio optimization and the random magnet problem*, Europhys. Lett. **59** (2002) 500.
- [15] V. Plerou, P. Gopikrishnan, B. Rosenow, L.A.N. Amaral, T. Guhr, and H.E. Stanley, *A Random Matrix Theory Approach to Cross-Correlations in Financial Data*, Phys. Rev. E **65** (2002) 066126.
- [16] S. Sharifi, M. Crane, A. Shamaie, and H. Ruskin, *Random Matrix Theory for Portfolio Optimization: A Stability Approach*, Physica A **335** (2004) 629-643.
- [17] T. Conlon, H.J. Ruskin, and M. Crane, *Random Matrix Theory and Fund of Funds Portfolio Optimisation*, Physica A **382** (2007) 565-578.
- [18] E. P. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions*, Ann. Math. **62** (1955), 548-564.
- [19] E. P. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. Math. **67** (1958), 325-327.
- [20] S. Pafka and I. Kondor, *Noisy covariance matrices and portfolio optimization*, Eur. Phys. J. B **27** (2002) 277-280.
- [21] J-P. Onnela, A. Chakraborti, and K. Kaski, *Dynamics of market correlations: taxonomy and portfolio analysis*, Phys. Rev. E **68** (2003) 056110.
- [22] V. Tola, F. Lillo, M. Gallegati, and R.N. Mantegna, *Cluster analysis for portfolio optimization*, Journal of Economic Dynamics and Control **32** (2008) 235-258.
- [23] E. Pantaleo, M. Tumminello, F. Lillo, and R.S. Mantegna, *When do improved covariance matrix estimators enhance portfolio optimization? An empirical comparative study of nine estimators*, Quantitative Finance **11** (2011) 1067-1080.

- [24] J-P Bouchaud and M. Potters, *Financial applications of random matrix theory: a short review*, to appear in the “Handbook on random matrix theory”, Oxford University Press.
- [25] V.A. Marenko and L.A. Pastur, USSR-Sb **1** (1967) 457-483.

A Stocks of the BM&F-Bovespa used for the portfolio building

2004-2005

AMBV4	BBAS3	BBDC3	BBDC4	BRAP4	BRKM5	BRTO4	CCRO3	CESP5	CGAS5
CLSC6	CMIG3	CMIG4	CNFB4	CPLE3	CPLE6	CRUZ3	CSNA3	CTNM4	EBTP3
EBTP4	ELET3	ELET6	EMBR3	FESA4	FFTL4	FIBR3	GGBR4	GOAU4	INEP4
ITSA4	ITUB4	KLBN4	LAME4	LIGT3	MAGG3	PCAR5	PETR3	PETR4	POMO4
RAPT4	SBSP3	SUZB5	TBLE3	TCSL3	TCSL4	TLPP3	TLPP4	TMAR5	TNCP3
TNCP4	TNLP3	TNLP4	TRPL4	UNIP6	USIM5	VALE3	VALE5	VIVO3	VIVO4
WEGE3									

2005-2006

AMBV4	BBAS3	BBDC3	BBDC4	BRAP4	BRKM5	BRTO4	CCRO3	CESP5	CGAS5
CLSC6	CMIG3	CMIG4	CNFB4	CPFE3	CPLE6	CRUZ3	CSNA3	CTNM4	DASA3
EBTP3	EBTP4	ELET3	ELET6	EMAE4	EMBR3	FFTL4	FIBR3	FJTA4	GGBR3
GGBR4	GOAU4	GOLL4	GRND3	IDNT3	ITSA4	ITUB4	KLBN4	LAME4	LIGT3
MAGG3	MGEL4	NATU3	PCAR5	PETR3	PETR4	POMO4	PSSA3	RAPT4	SBSP3
SUZB5	TBLE3	TCSL3	TCSL4	TELB3	TELB4	TLPP3	TLPP4	TMAR5	TNCP3
TNCP4	TNLP3	TNLP4	TRPL4	UGPA4	UNIP6	USIM5	VALE3	VALE5	VIVO3
VIVO4	WEGE3								

2006-2007

ALLL3	AMBV3	AMBV4	BBAS3	BBDC3	BBDC4	BOBR4	BRAP4	BRKM5	BRTO4
BTOW3	CCRO3	CGAS5	CLSC6	CMIG3	CMIG4	CNFB4	CPFE3	CPLE6	CRUZ3
CSAN3	CSNA3	CTAX4	CTNM4	CYRE3	DASA3	ELET3	ELET6	EMAE4	EMBR3
ENBR3	ENMA3B	ETER3	FFTL4	FIBR3	FJTA4	GGBR3	GGBR4	GOAU4	GOLL4
GRND3	GUAR3	IDNT3	ITSA4	ITUB4	KLBN4	LAME4	LIGT3	LREN3	MAGG3
MGEL4	NATU3	NETC4	OHLB3	PCAR5	PETR3	PETR4	POMO4	PSSA3	RAPT4
RENT3	RSID3	SBSP3	SLED4	SUZB5	TAMM4	TBLE3	TCSL3	TCSL4	TELB3
TELB4	TLPP3	TLPP4	TMAR5	TNLP3	TNLP4	TRPL4	UGPA4	UNIP6	UOLL4
USIM5	VALE3	VALE5	VIVO3	VIVO4	WEGE3				

2007-2008

ALLL3	AMBV3	AMBV4	BBAS3	BBDC3	BBDC4	BEES3	BISA3	BRAP4	BRFS3
BRKM5	BRTO4	BTOW3	CARD3	CCRO3	CESP6	CGAS5	CLSC6	CMIG3	CMIG4
CNFB4	COCE5	CPFE3	CPLE6	CRUZ3	CSAN3	CSMG3	CSNA3	CTNM4	CYRE3
DASA3	ECOD3	ELET3	ELET6	ELPL4	EMBR3	ENBR3	EQTL3	ESTR4	ETER3
FFTL4	FIBR3	FJTA4	GETI3	GETI4	GFS43	GGBR3	GGBR4	GOAU4	GOLL4
GRND3	IDNT3	IMBI4	ITSA4	ITUB4	JBDU4	KLBN4	LAME4	LIGT3	LPSB3
LREN3	LUPA3	MAGG3	MDIA3	MLFT4	NATU3	NETC4	ODPV3	OHLB3	PCAR5
PETR3	PETR4	POMO4	POS13	PSSA3	RAPT4	RENT3	RNAR3	RSID3	SBSP3
SLED4	SUZB5	TAMM4	TBLE3	TCSL3	TCSL4	TELB3	TELB4	TLPP3	TLPP4
TMAR5	TNLP3	TNLP4	TOTS3	TRPL4	UGPA4	UNIP6	UOLL4	USIM3	USIM5
VALE3	VALE5	VIVO4	VLID3	WEGE3					

2008-2009

ABCB4	ALLL3	AMAR3	AMBV3	AMBV4	AMIL3	BAZA3	BBAS3	BBDC3	BBDC4
BEEF3	BEES3	BEMA3	BICB4	BISA3	BMT04	BPNM4	BRAP4	BRFS3	BRKM5
BRML3	BRSR6	BRT04	BTOW3	BVMF3	CARD3	CCIM3	CCRO3	CESP6	CGAS5
CLSC6	CMIG3	CMIG4	CNFB4	COCE5	CPFE3	CPLE6	CRUZ3	CSAN3	CSMG3
CSNA3	CYRE3	CZRS4	DASA3	DAYC4	ECOD3	ELET3	ELET6	ELPL4	EMBR3
ENBR3	EQTL3	ESTR4	ETER3	EVEN3	EZTC3	FFTL4	FHER3	FJTA4	GETI3
GETI4	GFSA3	GGBR3	GGBR4	GOAU4	GOLL4	GPIV11	GRND3	IDNT3	IGTA3
INPR3	ITSA4	ITUB3	ITUB4	JBDU4	JBSS3	JHSF3	KEPL3	KLBN4	LAME4
LIGT3	LOGN3	LPSB3	LREN3	LUPA3	MAGG3	MILK11	MLFT4	MMXM3	MRFG3
MRVE3	MULT3	NATU3	NETC4	ODPV3	OHLB3	PCAR5	PDGR3	PETR3	PETR4
PINE4	PLAS3	PMAM3	POMO4	POSI3	PRVI3	PSSA3	RAPT4	RDCD3	RDN13
RENT3	ROMI3	RSID3	SBSP3	SFSA4	SLCE3	SLED4	SMT03	SULA11	SUZB5
TAMM4	TBLE3	TCSA3	TCSL3	TCSL4	TELB3	TELB4	TGMA3	TLPP3	TLPP4
TMAR5	TNLP3	TNLP4	TOTS3	TOYB3	TOYB4	TRPL4	UGPA4	UNIP6	UOLL4
USIM3	USIM5	VALE3	VALE5	VIVO4	VLID3	WEGE3	WSON11		

2009-2010

ABCB4	ALLL3	AMAR3	AMBV3	AMBV4	AMIL3	BBAS3	BBDC3	BBDC4	BEEF3
BEMA3	BICB4	BISA3	BPNM4	BRAP4	BRFS3	BRKM5	BRML3	BRSR6	BRT04
BTOW3	BVMF3	CARD3	CCIM3	CCRO3	CESP6	CLSC6	CMIG3	CMIG4	CNFB4
COCE5	CPFE3	CPLE6	CRDE3	CRUZ3	CSAN3	CSMG3	CSNA3	CYRE3	DASA3
ECOD3	ELET3	ELET6	ELPL4	EMBR3	ENBR3	EQTL3	ESTR4	ETER3	EVEN3
EZTC3	FESA4	FFTL4	FHER3	FJTA4	GETI3	GETI4	GFSA3	GGBR3	GGBR4
GOAU4	GOLL4	GPIV11	GRND3	GSHP3	HBOR3	HYPE3	IDNT3	IGTA3	INEP4
INPR3	ITSA4	ITUB3	ITUB4	JBDU4	JBSS3	JHSF3	KEPL3	KLBN4	KROT11
LAME3	LAME4	LIGT3	LLXL3	LOGN3	LPSB3	LREN3	LUPA3	MAGG3	MILK11
MLFT4	MMXM3	MNDL4	MPXE3	MRFG3	MRVE3	MULT3	MYPK3	NATU3	NETC4
ODPV3	OGXP3	OHLB3	PCAR5	PDGR3	PETR3	PETR4	PINE4	PLAS3	PMAM3
POMO4	POSI3	PRVI3	PSSA3	RAPT4	RDCD3	RENT3	RSID3	SBSP3	SFSA4
SLCE3	SLED4	SMT03	SULA11	SUZB5	TAMM4	TBLE3	TCSA3	TCSL3	TCSL4
TELB3	TELB4	TGMA3	TLPP3	TLPP4	TMAR5	TNLP3	TNLP4	TOTS3	TOYB3
TOYB4	TPIS3	TRPL4	UGPA4	UNIP6	UOLL4	USIM3	USIM5	VALE3	VALE5
VIVO4	VLID3	WEGE3							